arXiv:math/0411534v2 [math.NT] 25 Nov 2004

HEEGNER POINTS AND MORDELL-WEIL GROUPS OF ELLIPTIC CURVES OVER LARGE FIELDS

BO-HAE IM

Abstract. Let $E/\mathbb{Q}$ be an elliptic curve defined over $\mathbb{Q}$ with conductor N and $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ the absolute Galois group of an algebraic closure $\overline{\mathbb{Q}}$ of $\mathbb{Q}$. We prove that for every $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, the Mordell-Weil group $E(\overline{\mathbb{Q}}^{\sigma})$ of E over the fixed subfield of $\overline{\mathbb{Q}}$ under σ has infinite rank. Our approach uses the modularity of $E/\mathbb{Q}$ and a collection of algebraic points on E – the so-called *Heegner points* – arising from the theory of complex multiplication. In particular, we show that for some integer r, the rank of E over all the ring class fields of conductor of the form rm, and of the form rp^n is unbounded, as m goes to infinity, as m and as n goes to infinity respectively, where m is a square-free integer and p is a prime such that $(m, rN) = 1$ and $(p, rN) = 1$.

This paper is motivated by the following conjecture of M. Larsen [6]:

Conjecture. Let K be a number field and E/K an elliptic curve over K. Then, for every $\sigma \in \mathrm{Gal}(\overline{K}/K)$, the Mordell-Weil group $E(\overline{K}^{\sigma})$ of E over $\overline{K}^{\sigma} = \{x \in \overline{K} \mid \sigma(x) = x\}$ has infinite rank.

In [3], we have proved this conjecture in certain cases:
for a number field K and an elliptic curve E/K over K,

- if 2-torsion points of E/K are K-rational, or
- if E/K has a K-rational point P such that $2P \neq O$ and $3P \neq O$,

then for every automorphism $\sigma \in \mathrm{Gal}(\overline{K}/K)$, the rank of the Mordell-Weil group $E(\overline{K}^{\sigma})$ is infinite.

In this paper, we prove that the conjecture is true for elliptic curves over $\mathbb{Q}$ without any hypothesis on rational points of $E/\mathbb{Q}$, *i.e.* if $E/\mathbb{Q}$ is an elliptic curve over $\mathbb{Q}$, then,

---

Date: July 26, 2004.

2000 *Mathematics Subject Classification.* Primary 11G05.

for every automorphism $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, the Mordell-Weil group $E(\overline{\mathbb{Q}}^{\sigma})$ over the fixed subfield of $\overline{\mathbb{Q}}$ under σ has infinite rank.

To prove the conjecture for a given E/K, ultimately one must find an infinite supply of rational points of E over finite extensions of K contained in $\overline{K}^{\sigma}$. In [3], we constructed such points using Diophantine geometry, essentially by searching for sufficiently rational subvarieties of certain quotients of the n-fold product E^n of E.

Here we use a completely different approach, coming from arithmetic: taking advantage of the modularity of elliptic curves over $\mathbb{Q}$, we choose our rational points on E to be algebraic points over ring class fields – the so-called *Heegner points.* The main strategy is as follows: by using the norm compatibility properties of Heegner points and a generalized dihedral group structure of the Galois groups of ring class fields over $\mathbb{Q}$, we show that the rank of E over the ring class fields is unbounded as the ring class fields get larger. And we also show that a given automorphism $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ does not fix a quadratic imaginary extension of $\mathbb{Q}$ over which all primes dividing the conductor of E are split, then the rank of E over the fixed subfields of the ring class fields under σ is unbounded as the ring class fields get larger, which proves that the rank of E over $\overline{\mathbb{Q}}^{\sigma}$ is infinite. On the other hand, if $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ fixes infinitely many quadratic imaginary extensions, then we can construct infinitely many linearly independent points defined over each of them by applying the Hilbert irreducibility theorem to a Weierstrass equation of $E/\mathbb{Q}$ directly and this also proves infinite rank of E over $\overline{\mathbb{Q}}^{\sigma}$.

Acknowledgements

I would like to thank my thesis advisor, Michael Larsen for suggesting this problem and for valuable discussions and helpful comments on this paper. Also, I wish to thank Henri Darmon for suggesting this approach, and for his guidance and valuable advice and comments on an earlier manuscript of this paper.

1. THE MAIN THEOREM

In this section, we introduce the main theorem. First, we will need the following Hilbert irreducibility and the denseness of Hilbert sets in any open intervals of $\mathbb{R}$.

Let $f \in K(t_1, \dots, t_m)[X_1, \dots, X_n]$ be a polynomial with coefficients in the quotient field $K(t_1, \dots, t_m)$ of $K[t_1, \dots, t_m]$ which is irreducible over $K(t_1, \dots, t_m)$. We define

$$H_K(f) = \{(a_1, \dots, a_m) \in K^m \mid f(a_1, \dots, a_m, X_1, \dots, X_n) \text{ is irreducible over } K\}$$

to be the Hilbert set of f over K. If for every $m \geq 1$, any intersections of a finite number of Hilbert sets with a finite number of nonempty Zariski open subsets in K^m are not empty (in fact, they are infinite), a field K is called *a Hilbertian field*.

Lemma 1.1. *Let L be a finite separable extension of a Hilbertian field K and let f be a polynomial in $L(t_1, \dots, t_m)[X_1, \dots, X_n]$ which is irreducible over the quotient field $L(t_1, \dots, t_m)$. Then, there exists a polynomial $p \in K[t_1, \dots, t_m, X_1, \dots, X_n]$ such that p is irreducible over $K(t_1, \dots, t_m)$ and $H_K(p) \subseteq H_L(f)$.*

Proof. For a given irreducible polynomial $f \in L(t_1, \dots, t_m)[X_1, \dots, X_n]$, by ([4], Ch.11, Lemma 11.6), there is an irreducible polynomial $q \in K(t_1, \dots, t_m)[X_1, \dots, X_n]$ such that $H_K(q) \subseteq H_L(f)$. By ([4], Ch.11, Lemma 11.1), there is an irreducible polynomial $p \in K[t_1, \dots, t_m, X_1, \dots, X_n]$ which is irreducible over $K(t_1, \dots, t_m)$ such that $H_K(p) \subseteq H_K(q)$. Hence the Hilbert set $H_L(f)$ of f over L contains the Hilbert set $H_K(p)$ of p over K. □

Lemma 1.2. *Let K be a number field and $\tau_1, \dots, \tau_m$ be a family of real embeddings of K. For $i = 1, 2, \dots, k$, let $f_i(x, y) \in K[x, y]$ be irreducible polynomials over $K(x)$. Let $H_K(f_i) = \{\alpha \in K \mid f_i(\alpha, y) \in K[y]$ is irreducible over $K\}$ be the Hilbert set of f_i over K. Then*

$$\left(\bigcap_{i=1}^{k} H_K(f_i)\right) \cap \left(\bigcap_{j=1}^{m} \tau_j^{-1}(I)\right) \neq \emptyset,$$

for any open interval I in $\mathbb{R}$.

Proof. Since K is a finite separable extension of $\mathbb{Q}$, by Lemma 1.1, there exist irreducible polynomials $F_i(x, y) \in \mathbb{Q}[x, y]$ such that for each $i = 1, 2, \dots, k$, the Hilbert set $H_{\mathbb{Q}}(F_i)$ of F_i over $\mathbb{Q}$ is contained in the Hilbert set $H_K(f_i)$ of f_i over K.

Let I be an open interval in $\mathbb{R}$. Since $\bigcap_{i=1}^{k} H_{\mathbb{Q}}(F_i)$ is dense in $\mathbb{Q}$ by ([5], Chapter 9, Corollary 2.5), and $\mathbb{Q}$ is dense in $\mathbb{R}$, $\left(\bigcap_{i=1}^{k} H_{\mathbb{Q}}(F_i)\right) \cap I$ is not empty. Hence there is

$\beta \in \left(\bigcap_{i=1}^{k} H_{\mathbb{Q}}(F_i)\right) \cap I$. Since $\bigcap_{i=1}^{k} H_{\mathbb{Q}}(F_i) \subseteq \bigcap_{i=1}^{k} H_K(f_i)$, we have $\beta \in \bigcap_{i=1}^{k} H_K(f_i)$. On the other hand, for each real embedding τ_j of K, we have $\tau_j|_{\mathbb{Q}} = id_{\mathbb{Q}}$. Hence for all $j = 1, 2, \dots, m$, $\tau_j(\beta) = \beta \in I$. Hence $\beta \in \bigcap_{j=1}^{m} \tau_j^{-1}(I)$. Therefore, $\beta \in \left(\bigcap_{i=1}^{k} H_K(f_i)\right) \cap \left(\bigcap_{j=1}^{m} \tau_j^{-1}(I)\right)$. □

Here is our main theorem.

Theorem 1.3. *Let $E/\mathbb{Q}$ be an elliptic curve over $\mathbb{Q}$. Then, for every automorphism $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, the rank of $E(\overline{\mathbb{Q}}^{\sigma})$ is infinite.*

Proof. Let N be the conductor of E and let $y^2 = x^3 + ax + b$ be a Weierstrass equation of $E/\mathbb{Q}$. By the change of variables, we may assume that a and b are integers. Let $M = 4p_1p_2 \cdots p_k$, where p_k are all distinct prime factors of N. Consider the polynomial

$$f(x) = (1 + Mx)^3 + aM^4(1 + Mx) + bM^6 \ \in \mathbb{Z}[x].$$

Then, there exists a real number r such that for all $x < r$, the expression $f(r)$ is strictly negative. Let $I = (-\infty, r)$ be the open interval in $\mathbb{R}$ of all real numbers less than r. Since $\mathbb{Q}$ is Hilbertian, by Lemma 1.2, there exists an integer $m_1 \in I$ such that $y^2 - f(m_1)$ is irreducible over $\mathbb{Q}$. Let $K_{m_1} = \mathbb{Q}(\sqrt{f(m_1)})$ be the quadratic imaginary extension of $\mathbb{Q}$. By Lemma 1.1, there is a polynomial p over $\mathbb{Q}$ such that $H_{\mathbb{Q}}(p) \subseteq H_{K_{m_1}}(y^2 - f(x))$. Then, by Lemma 1.2 again, there exists an integer $m_2 \in I \cap H_{\mathbb{Q}}(p)$. So $K_{m_2} = \mathbb{Q}(\sqrt{f(m_2)})$ is a quadratic imaginary extension of $\mathbb{Q}$ and K_{m_1} and K_{m_2} are distinct, hence linearly disjoint over $\mathbb{Q}$. By repeating this procedure over the composite field of quadratic imaginary extensions obtained from the previous steps inductively, there is an infinite set S of integers such that for all $m \in S$

1. $f(m) < 0$, so that $K_m := \mathbb{Q}(\sqrt{f(m)})$ is a quadratic imaginary extension of $\mathbb{Q}$,
2. the fields in the infinite sequence $\{K_m\}_{m \in S}$ are linearly disjoint over $\mathbb{Q}$ and
3. if $E/\mathbb{Q}$ has CM, then, K_m is different from $F = \mathrm{End}(E) \otimes \mathbb{Q}$.

Note that for each $m \in S$ and for every prime p_i dividing N,

$$f(m) \equiv \begin{cases} 1 \pmod{p_i}, & \text{if } p_i \neq 2, \\ 1 \pmod{8}, & \text{if } p_i = 2. \end{cases}$$

Hence, this implies that all primes dividing N are split in K_m. On the other hand, the discriminant of K_m is $f(m)$ or $4f(m)$ depending on whether $f(m) \equiv 1 \pmod 4$ or not, respectively. And in any case, the discriminant of K_m is prime to N.

Let $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. We divide into two cases. For the first case, suppose that for all $m \in S$, $\sigma|_{K_m} = id_{K_m}$. Then, for each $m \in S$, consider the number $\dfrac{1+Mm}{M^2} \in \mathbb{Q}$. By plugging this number into the given Weierstrass equation of $E/\mathbb{Q}$, we get

$$y^2 = \left(\frac{1+Mm}{M^2}\right)^3 + a\left(\frac{1+Mm}{M^2}\right) + b = \frac{f(m)}{M^6}.$$

Hence the point

$$P_m = \left(\frac{1+Mm}{M^2}, \frac{\sqrt{f(m)}}{M^3}\right)$$

is in $E(K_m)$ but it is not in $E(\mathbb{Q})$. And moreover, since $K_m = K_m^\sigma$, P_m is fixed under σ.

So we get an infinite sequence $\{P_m\}_{m\in S}$ of points in $E(\overline{\mathbb{Q}}^\sigma)$ such that each P_m is defined over the quadratic imaginary extension K_m over $\mathbb{Q}$. We may assume that these points are not torsion points by ([9], Lemma). Now we show the points P_m for $m \in S$ are linearly independent. Suppose that for some integers a_i and $m_i \in S$,

$$a_1 P_{m_1} + a_2 P_{m_2} + \cdots + a_k P_{m_k} = O.$$

Since the fields in $\{K_m\}_{m\in S}$ are linearly disjoint over $\mathbb{Q}$, for each i, there is an automorphism of $\overline{\mathbb{Q}}$ which fixes all but one K_{m_i} of $K_{m_1}, \ldots, K_{m_k}$. Note that such an automorphism takes P_{m_i} to its inverse, $-P_{m_i}$. Applying this automorphism, we get

$$a_1 P_{m_1} + \cdots + a_{i-1} P_{m_{i-1}} - a_i P_{m_i} + \cdots + a_k P_{m_k} = O.$$

By subtracting, we get $2a_i P_{m_i} = O$, which implies $a_i = 0$. We conclude that the P_m for $m \in S$ are linearly independent in $E(\overline{\mathbb{Q}}^\sigma) \otimes \mathbb{Q}$. Hence the rank of $E(\overline{\mathbb{Q}}^\sigma)$ is infinite.

For the second case, suppose that there is an integer $m \in S$ such that $\sigma|_{K_m} \neq id_{K_m}$. Then, fix such a quadratic imaginary extension K_m, and call it K, and let K_{ab} be the maximal abelian extension of K. Then, we complete the proof of this case as a consequence of the following stronger statement:

Theorem 1.4. *Under the assumption in the second case above (i.e. if K is different from* $\mathrm{End}(E) \otimes \mathbb{Q}$*, all primes dividing N are split in K, and $\sigma|_K \neq id_K$), the rank of the Mordell-Weil group of E over the fixed subfield $(K_{ab})^\sigma$ of K_{ab} under σ is infinite.*

The proof of Theorem 1.4, which lies deeper than the methods used in the first case will be treated in the following section and will be given explicitly in Proposition 2.9, as it requires modularity of $E/\mathbb{Q}$ and the theory of complex multiplication which give a non-torsion algebraic point of E under the given assumption in the second case. $\square$

2. The rank of E over ring class fields of imaginary quadratic fields

The goal of this section is to prove Theorem 1.4 — stated at the end of the previous section. By a theorem of Wiles [12] and Taylor-Wiles [11] (completed by a later work of Breuil, Conrad, Diamond and Taylor [1]), the elliptic curve $E/\mathbb{Q}$ is known to be *modular*. Our strategy is to construct algebraic points on $E((K_{ab})^{\sigma})$ using *Heegner points* over the ring class fields arising from the theory of complex multiplication.

We will need the following lemma later.

Lemma 2.1. *Let K be an imaginary quadratic extension of $\mathbb{Q}$ such that K is different from $F = \mathrm{End}(E) \otimes \mathbb{Q}$. For a prime ℓ, let a_l be the coefficient of the Hecke operator T_ℓ of the modular form of E. Then, there is an integer M such that for all $p \geq M$ (which is inert in F, if E has CM), there is a prime q such that*

1. *q is inert in K,*
2. *p does not divide a_q and*
3. *p divides $q+1$.*

Proof. Suppose $E/\mathbb{Q}$ has no CM. Then, by ([8], §4. Theorem 2), there is a large integer M such that for all primes $p \geq M$, the continuous Galois representation

$$\rho_p : \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \to \mathbf{GL}_2(\mathbb{F}_p)$$

is surjective. In particular, let M be large enough such that every $p \geq M$ is unramified in K, since K is ramified at only finitely many primes.

Since $Ker(\rho_p)$ is an open normal subgroup of $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, there is a finite extension L over $\mathbb{Q}$ such that $Ker(\rho_p) = \mathrm{Gal}(\overline{\mathbb{Q}}/L)$.

And since

$$\mathrm{Gal}(L/\mathbb{Q}) \cong \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})/Ker(\rho_p) \cong \mathbf{GL}_2(\mathbb{F}_p)$$

has a unique subgroup of index 2, the kernel of $det^{\frac{p-1}{2}}$, the unique quadratic field L' over $\mathbb{Q}$ in L is ramified only at p. On the other hand, since p is unramified in K, the fields

K and L' are linearly disjoint. Let S be a finite set of primes such that the composite field KL' is unramified and K is unramified outside S. Then, by the Cebotarev density theorem, each Frobenius automorphism Frob_q comes up infinitely often in $\mathrm{Gal}(KL'/\mathbb{Q})$, for $q \notin S$.

Since K and L' are linearly disjoint, we can choose the Frobenius automorphism Frob_q in $\mathrm{Gal}(K/\mathbb{Q})$ and $\mathrm{Gal}(L'/\mathbb{Q})$, independently. Hence, there is a prime $q \neq 2$ such that q is inert in K and

$$\rho_p(\mathrm{Frob}_q) = \begin{pmatrix} a & b \\ 0 & -a^{-1} \end{pmatrix}$$

where a is in $\mathbb{F}_p$ such that the order of a is not 2 in $\mathbb{F}_p$ and b is some element of $\mathbb{F}_p$. Then, we have

$$a_q = \mathrm{Trace}(\rho_p(\mathrm{Frob}_q)) = a - a^{-1} \not\equiv 0 \pmod{p}.$$

This implies that p does not divide a_q, which is condition (2). Also, we have

$$q = \det(\rho_p(\mathrm{Frob}_q)) \equiv -1 \pmod{p}.$$

This implies that p divides $q+1$ which is condition (3). Hence such a prime q satisfies all conditions (1) though (3).

Suppose $E/\mathbb{Q}$ has CM. Let $R = \mathrm{End}(E)$ and $F = R \otimes \mathbb{Q}$. Then, F is an imaginary quadratic extension of $\mathbb{Q}$ such that $R = \mathrm{End}_F(E)$.

By ([8], §4. Corollary of Theorem 5), there is a large integer M such that for all primes $p \geq M$, the continuous Galois representation ρ_p factors through a surjective homomorphism,

$$\sigma_p : \mathrm{Gal}(\overline{F}/F) \to R_p^*,$$

where $R_p = R \otimes \mathbb{Z}_p \cong \mathbb{Z}_p^2$ as $\mathbb{Z}_p$-modules. Under the canonical embedding $R_p = \mathrm{End}_{R_p}(R_p) \subseteq \mathrm{End}_{\mathbb{Z}_p}(\mathbb{Z}_p^2) = M_2(\mathbb{Z}_p)$, the norm map and the trace map from R_p to $\mathbb{Z}_p$ are the restrictions of the determinant map and the trace map on $M_2(\mathbb{Z}_p)$.

Let $\mathcal{O}$ be the maximal order in F. Then, $\mathcal{O}/R$ is finite, so if $p \gg 0$,

$$R/pR \cong \mathcal{O}/p\mathcal{O} \cong \mathbb{F}_p \times \mathbb{F}_p \text{ or } \mathbb{F}_{p^2},$$

depending on whether p is split or inert in F. In either case, the norm map N is a non-degenerate quadratic form on the underlying 2-dimensional vector space over $\mathbb{F}_p$. For any non-zero k, $N(\alpha) - kz^2$ is a non-degenerate quadratic form on the 3-dimensional $\mathbb{F}_p$-vector space $R/pR \times \mathbb{F}_p$. In particular, $N(\alpha) = -z^2$ defines a conic curve. Since any conic has a rational point over a finite field by the Chevalley-Warning theorem ([7], Chapter I, Theorem 3), the conic $N(\alpha) = -z^2$ over $\mathbb{F}_p$ is isomorphic to $\mathbb{P}^1$, hence it has $p+1$ points. Since $N(\alpha) = 0$ (when $z = 0$) has at most two non-trivial solutions on the line at ∞, there are at least $p+1-3$ points of $N(\alpha) = -1$. Therefore, for any prime $p \geq 5$, $N(\alpha) = -1$ has at least three solutions over $\mathbb{F}_p$. On the other hand, the trace map T is a linear form on the underlying 2-dimensional vector space. And the system of two equations, $T(\alpha) = 0$ and $N(\alpha) = -1$ has at most two solutions. Since $N(\alpha) = -1$ has at least three solutions for any prime $p \geq 5$, we can find $\alpha \in R/pR$ such that

$$T(\alpha) \not\equiv 0 \text{ and } N(\alpha) \equiv -1 \pmod{p}.$$

Since the norm $N(\alpha)$ of α is congruent to -1 (mod p), we can lift α to a p-adic unit $\widetilde{\alpha} \in R_p^*$ by applying the Hensel's lemma to the norm map.

Let $M \geq 5$ be large enough that for all $p \geq M$, $R/pR \cong \mathcal{O}/p\mathcal{O}$ and let $p \geq M$ be inert in F and unramified in K.

Let d_p denote the reduction of $det(\rho_p)$ mod p, and let $\beta_p = d_p^{\frac{p-1}{2}}$. Then, $Ker(\beta_p)$ is of index 2 in $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ and therefore, corresponds to a quadratic extension L which is ramified only at p. Thus, the fields, F, K and L are all linearly disjoint over $\mathbb{Q}$. In particular, $\sigma_p(\mathrm{Gal}(\overline{F}/FL))$ is of index 2 in R_p^*, which has a unique open subgroup of index 2, since p is inert in F. As $FK \neq FL$, K is not contained in the field associated to $Ker(\sigma_p)$, which implies K is linearly disjoint from this field. By the Cebotarev density theorem, there exists q which is inert in K and split in F and such that

$$\sigma_p(\mathrm{Frob}_q) = \widetilde{\alpha} \in R_p^*.$$

Then, we have

$$a_q = T(\sigma_p(\mathrm{Frob}_q)) = T(\widetilde{\alpha}) \not\equiv 0 \pmod{p}.$$

This implies that p does not divide a_q, which is condition (2). Also, we have

$$q = \det(\sigma_p(\mathrm{Frob}_q)) = N(\widetilde{\alpha}) \equiv -1 \pmod{p}.$$

This implies that p divides $q+1$ which is condition (3). This completes the proof. $\square$

For a given elliptic curve $E/\mathbb{Q}$, we fix an element $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ and a quadratic imaginary extension K of $\mathbb{Q}$ which is not fixed under σ. And we assume that all primes dividing the conductor N of $E/\mathbb{Q}$ are split in K. Note that this setting corresponds to the second case in the proof of Theorem 1.3.

For each integer n relatively prime to N, let H_n denote the ring class field of K of conductor n. By ([2], Chapter 3, Theorem 3.6), there is an algebraic point $P_n \in E(H_n)$ which is called a *Heegner point of conductor* n. Let $HP(n) \subset E(H_n)$ denote the set of all Heegner points of conductor n in $E(H_n)$. Then, the set $HP(n)$ satisfies the following properties.

First, we recall the norm-compatibility properties of Heegner points.

Proposition 2.2. *Let $E/\mathbb{Q}$ be a modular elliptic curve over $\mathbb{Q}$ and N the conductor of $E/\mathbb{Q}$. Let n be an integer and ℓ a prime number such that both n and ℓ are prime to N. Let $P_{n\ell}$ be any point in $HP(n\ell)$ and a_ℓ the coefficient of the Hecke operator T_ℓ of the modular form of E. Then, there are points $P_n \in HP(n)$ and (when $\ell|n$) $P_{n/\ell} \in HP(n/\ell)$ such that*

$$Trace_{H_{nl}/H_n}(P_{nl}) = \begin{cases} a_l P_n, & \text{if } l \nmid n \text{ is inert in } K \\ (a_l - \sigma_\lambda - \sigma_\lambda^{-1})P_n, & \text{if } l = \lambda\bar{\lambda} \nmid n \text{ is split in } K \\ (a_l - \sigma_\lambda)P_n, & \text{if } l = \lambda^2 \text{ is ramified in } K \\ a_l P_n - P_{n/l}, & \text{if } l|n. \end{cases}$$

Proof. See ([2], Chapter 3, Proposition 3.10). ☐

Lemma 2.3. *Let H_∞ be the union of all the ring class fields of conductor prime to N. Then, the set $E(H_\infty)_{tor}$ is finite.*

Proof. See ([2], Chapter 3, Lemma 3.14). ☐

The following lemma describes the structure of the Galois groups of ring class field over an imaginary quadratic extension K of $\mathbb{Q}$.

Lemma 2.4. *Let N be the conductor of $E/\mathbb{Q}$. Let H_n be the ring class field of conductor n over an imaginary quadratic extension K over $\mathbb{Q}$.*

(a) If p is a prime not dividing $c \cdot N \cdot [H_c : K] \cdot disc(H_c)$, then for all integers $n \geq 1$,

$$\mathrm{Gal}(H_{cp^n}/H_{cp}) \cong \mathbb{Z}/p^{n-1}\mathbb{Z}, \text{ and } \mathrm{Gal}(H_{cp^{n+1}}/H_{cp^n}) \cong \mathbb{Z}/p\mathbb{Z}.$$

(b) If $k = \prod_{j=1}^{m} p_j$ *for distinct primes* p_j *which are relatively prime to* N *and inert in* K*, then for each* j*,*

$$\mathrm{Gal}(H_k/H_{\frac{k}{p_j}}) \cong \mathbb{Z}/(p_j+1)\mathbb{Z}.$$

Proof. To prove the first part of (a), for $n \geq 1$ and $k \geq 1$,

$$\begin{aligned}\mathrm{Gal}(H_{cp^n}/H_{cp}) &\cong ker(\mathrm{Gal}(H_{cp^n}/K) \to \mathrm{Gal}(H_{cp}/K)) \\ &\cong ker((\mathcal{O}_K/cp^n\mathcal{O}_K)^*/(\mathbb{Z}/cp^n\mathbb{Z})^* \to (\mathcal{O}_K/cp\mathcal{O}_K)^*/(\mathbb{Z}/cp\mathbb{Z})^*) \\ &= \frac{[(1+cp\mathcal{O}_K)/cp^n\mathcal{O}_K]^* \cdot (\mathbb{Z}/cp^n\mathbb{Z})^*}{(\mathbb{Z}/cp^n\mathbb{Z})^*} \\ &= \frac{[(1+cp\mathcal{O}_K)/cp^n\mathcal{O}_K]^*}{[(1+cp\mathbb{Z})/cp^n\mathbb{Z}]^*} \\ &= \frac{[(1+p\mathcal{O}_K)/p^n\mathcal{O}_K]^*}{[(1+p\mathbb{Z})/p^n\mathbb{Z}]^*}, \text{ (since } p \nmid c \cdot [H_c : K] \cdot disc(H_c).) \\ &\cong \frac{[(1+p\mathcal{O}_K)/p^n\mathcal{O}_K]^*}{(\mathbb{Z}/p^{n-1}\mathbb{Z})}, \\ &\quad \text{(since } [(1+p\mathbb{Z})/p^n\mathbb{Z}]^* \cong ker((\mathbb{Z}/p^n\mathbb{Z})^* \to (\mathbb{Z}/p\mathbb{Z})^*)) \cong \mathbb{Z}/p^{n-1}\mathbb{Z}.)\end{aligned}$$

By the logarithmic function from $1+p\widehat{\mathcal{O}}_{K,p} \to p\widehat{\mathcal{O}}_{K,p}$ which maps $1+p^n\widehat{\mathcal{O}}_{K,p} \mapsto p^n\widehat{\mathcal{O}}_{K,p}$, where $\widehat{\mathcal{O}}_{K,p}$ is the completion of $\mathcal{O}_K$ at p,

$$[(1+p\widehat{\mathcal{O}}_{K,p})/p^{n-1}\widehat{\mathcal{O}}_{K,p}]^* \cong \widehat{\mathcal{O}}_{K,p}/p^{n-1}\widehat{\mathcal{O}}_{K,p} \cong \mathcal{O}_K/p^{n-1}\mathcal{O}_K.$$

Hence, we have

$$\mathrm{Gal}(H_{cp^n}/H_{cp}) \cong (\mathcal{O}_K/p^{n-1}\mathcal{O}_K)/(\mathbb{Z}/p^{n-1}\mathbb{Z}) \cong \mathbb{Z}/p^{n-1}\mathbb{Z}.$$

For the second part of (a),

$$\begin{aligned}\mathrm{Gal}(H_{cp^{n+1}}/H_{cp^n}) &\cong ker(\mathrm{Gal}(H_{cp^{n+1}}/H_{cp}) \to \mathrm{Gal}(H_{cp^n}/H_{cp})) \\ &\cong ker((\mathbb{Z}/p^n\mathbb{Z})^* \to (\mathbb{Z}/p^{n-1}\mathbb{Z})^*) \\ &\cong \mathbb{Z}/p\mathbb{Z}.\end{aligned}$$

To prove (b), for each $i = 1, ..., m$,

$$\begin{aligned}\mathrm{Gal}(H_k/H_{\frac{k}{p_i}}) &\cong ker(\mathrm{Gal}(H_k/K) \to \mathrm{Gal}(H_{\frac{k}{p_i}}/K)) \\ &\cong ker((\mathcal{O}_K/k\mathcal{O}_K)^*/(\mathbb{Z}/k\mathbb{Z})^* \to (\mathcal{O}_K/\tfrac{k}{p_i}\mathcal{O}_K)^*/(\mathbb{Z}/\tfrac{k}{p_i}\mathbb{Z})^*) \\ &\cong ker\left(\prod_{j=1}^{m} \mathrm{Gal}(H_k/H_{\frac{k}{p_j}}) \to \prod_{j=1}^{i-1} \mathrm{Gal}(H_k/H_{\frac{k}{p_j}}) \times \{1\} \times \prod_{j=i+1}^{m} \mathrm{Gal}(H_k/H_{\frac{k}{p_j}})\right) \\ &\cong \mathrm{Gal}(H_k/H_{\frac{k}{p_i}}) \cong (\mathbb{Z}/\lambda\mathbb{Z})^*/(\mathbb{Z}/p_i\mathbb{Z})^*, \text{ where } \lambda \text{ is a prime factor of } p. \\ &\cong \mathbb{Z}/(p_i+1)\mathbb{Z}.\end{aligned}$$

□

A Heegner system attached to (E, K) is a collection of points $P_n \in E(H_n)$ indexed by integers n prime to the conductor N of E, and satisfying the norm compatibility properties given in ([2], Chapter 3, Proposition 3.10 including Proposition 2.2) and the behavior under the action of reflections described in ([2], Chapter 3, Proposition 3.11).

In our setting, since all primes dividing N are split in K (recall the construction of K in the proof of Theorem 1.3), there is a Heegner system in which at least one of the points P_n for some n is non-torsion by ([2], Chapter 3, Theorem 3.13). We call such a Heegner system a *non-trivial Heegner system*. We will need the following lemma.

Lemma 2.5. *Suppose there is a nontrivial Heegner system attached to* (E, K)*. Let* n *be a positive integer prime to the conductor of* E *such that there are non-torsion points of* $E(H_\infty)$ *which are not in* $E(H_n)$*. Then, there exists a positive integer* M *such that for all non-torsion points* $Q \in E(H_\infty)$ *such that* $Q \notin E(H_n)$*,* $mQ \notin E(H_n)$ *for all integers* $m > M$*.*

Proof. Suppose not. Then, for all M, there exist an integer $m > M$ and a non-torsion point $Q \in E(H_\infty) - E(H_n)$ such that $Q, 2Q, \ldots, (m-1)Q \notin E(H_n)$ but $mQ \in E(H_n)$.

We may assume that $m = p^k$ for some prime p dividing m and for a positive integer k by replacing Q by $\frac{m}{p^k}Q$. Then, either the exponent k or the prime p must go to infinity as M goes to infinity, since $m = p^k > M$.

Since $p^{k-1} < m$, $p^{k-1}Q \notin E(H_n)$. Hence there is an automorphism $\tau \in \mathrm{Gal}(H_\infty/H_n)$ such that $\tau(p^{k-1}Q) \neq p^{k-1}Q$. Hence for the nontrivial point $\tau(p^{k-1}Q) - p^{k-1}Q$,

$$p(\tau(p^{k-1}Q) - p^{k-1}Q) = \tau(p^kQ) - p^kQ = p^kQ - p^kQ = O.$$

Since p is a prime, this implies that the point $\tau(p^{k-1}Q) - p^{k-1}Q$ has order exactly p, hence the nontrivial point $\tau(Q) - Q$ has order exactly $m = p^k$. So we have shown that there are torsion points of order exactly m for all integers $m > M$. Therefore, as M goes to infinity, the order of torsion points in $E(H_\infty)$ is unbounded, which is a contradiction to the finiteness of the set of torsion points of $E(H_\infty)$ in Lemma 2.3. This completes the proof. □

Now, by using Lemma 2.1 and the norm-compatibility properties in Proposition 2.2, we prove that the subgroup generated by all of Heegner points of the given nontrivial Heegner system is not finitely generated as a subgroup of the elliptic curve over the union of all the ring class fields of conductor of the form rm for some r, where m is a square-free integer relatively prime to rN. Then, by using Lemma 2.5, we show the following unboundedness of the rank of Mordell-Weil groups over all of those ring class fields.

Proposition 2.6. *Let N be the conductor of the given elliptic curve $E/\mathbb{Q}$. If there is a nontrivial Heegner system attached to (E, K), where K is a quadratic imaginary extension of $\mathbb{Q}$ which is different from the field $\mathrm{End}(E) \otimes \mathbb{Q}$ such that there is a non-torsion Heegner point $P_r \in E(H_r)$ over some ring class field H_r of conductor r, then the rank of $E(H_{rm})$ is unbounded, as a square-free integer m such that $(m, rN) = 1$ goes to infinity.*

Proof. First, we show that the group generated by all of Heegner points of conductor rm for all square-free integers m such that $(m, rN) = 1$ in the given nontrivial Heegner system is not finitely generated. Suppose this group is finitely generated. Then, since there are only finitely many torsion (Heegner) points over all the ring class fields in the system by Lemma 2.3, for some integer n which is a square-free multiple of r and $(n/r, rN) = 1$, there is a fixed ring class field H_n over which all Heegner points of conductor rm for all square-free integers m are defined. And we may assume that a Heegner point P_n of level n is of infinite order by the assumption that there exists a non-torsion Heegner point $P_r \in E(H_r)$.

Only finitely many primes divide either P_n in $E(H_n)$ or points of $E(H_\infty)_{tor}$, because $E(H_n)$ is finitely generated by the Mordell-Weil Theorem and there are only finitely many torsion points over all the ring class fields in the system by Lemma 2.3. Let S be the finite set of primes which divide either P_n in $E(H_n)$ or the order of any point of the finite set $E(H_\infty)_{tor}$. Then, we can choose a large odd prime $p \notin S$ such that $p \geq n+2$ and if E has no CM, then p is unramified in K and if E has CM, then p is inert in the imaginary quadratic extension, $\mathrm{End}(E) \otimes \mathbb{Q}$ of $\mathbb{Q}$.

Then by Lemma 2.1, there is a prime q such that

(1) q is inert in K,
(2) p divides $q + 1$ and
(3) p does not divide a_q.

And since $q+1 \geq p$ by (2) and $p \geq n+2$, q is strictly greater than n. This implies q does not divide n. Therefore, the ring class field H_{nq} is a proper finite extension of H_n and nq is again a square-free multiple of r and $(nq/r, rN) = 1$.

By the norm-compatibility property given in Proposition 2.2, when $q \nmid n$ is inert in K, we have

$$\mathrm{Trace}_{H_{nq}/H_n}(P_{nq}) = a_q P_n.$$

On the other hand, $E(H_{nq}) = E(H_n)$, by assumption. Hence the trace of P_{nq} from H_{nq} to H_n is divisible by the degree of H_{nq} over H_n which is $q+1$ by (b) of Lemma 2.4.

Hence, $\mathrm{Trace}_{H_{nq}/H_n}(P_{nq})$ is divisible by p by the property (3). But by the property (4) and since $p \notin S$, p divides neither a_q nor the point P_n, which is a contradiction. So we have shown that the group generated by the Heegner points of conductor rm for all square-free integers m is not finitely generated. In particular, this shows that there is a non-torsion point $P_{nq} \in E(H_{nq})$ but not in $E(H_n)$.

By Lemma 2.5, for such a non-torsion point $P_{nq} \notin E(H_n)$, there exists an integer M such that $mP_{nq} \notin E(H_n)$ for all $m > M$. In other words, the point P_{nq} is independent of any points in $E(H_n)$. Hence, $E(H_n) \otimes \mathbb{Q} \neq E(H_{nq}) \otimes \mathbb{Q}$. Therefore, we conclude that the rank of $E(H_{rm})$ cannot be bounded, as a square-free integer m such that $(m, N) = 1$ goes to infinity. □

Now we prove the unboundedness of the rank of Mordell-Weil groups over all the ring class fields of conductor cp^n as n goes to infinity and this will be an important role in proving the main theorem. To prove this, we need the following simple lemma.

Lemma 2.7. *For an elliptic curve $E/\mathbb{Q}$ and for a prime p not dividing the conductor of E, let $a_p = p + 1 - \#E(\mathbb{F}_p)$. Then, there is no infinite sequence $\{c_n\}_{n=0}^{\infty}$ of integers with $c_0 \neq 0$ and satisfying the following linear recurrence,*

$$pc_{n+1} = a_p c_n - c_{n-1}, \text{ for } n \geq 1,$$

and for every N, there exists $n > N$ such that $c_n \neq 0$.

Proof. Suppose there is such an infinite sequence $\{c_n\}_{n=0}^{\infty}$ of integers satisfying the above conditions. Then, the linear recurrence implies that

$$(*) \qquad c_n = \alpha^n b_0 + \beta^n b_1, \text{ for all } n \geq 1,$$

where α and β are two solutions of the quadratic equation $x^2 - \frac{a_p}{p}x + \frac{1}{p} = 0$ and

$$b_0 = \left(\frac{-\beta}{\alpha-\beta}c_0\frac{1}{\alpha-\beta}c_1\right) \text{ and } b_1 = \left(\frac{\alpha}{\alpha-\beta}c_0 - \frac{1}{\alpha-\beta}c_1\right).$$

Note that this quadratic equation has no rational solutions, because if there was, the only possible pairs of rational solutions are 1 and $\frac{1}{p}$ or -1 and $-\frac{1}{p}$ and in either case, we get $a_p = \pm(p+1)$ which is impossible since $|a_p| < 2\sqrt{p}$ by Hesse's inequality ([10], Chapter V, Theorem 1.1). And $b_0 \neq 0$ and $b_1 \neq 0$ since c_0 is a nonzero integer and α and β are not rational numbers.

Let F be a quadratic extension of $\mathbb{Q}$ containing α and β and choose an embedding of F into $\overline{\mathbb{Q}}_p$ with the valuation v_p such that $v_p(\alpha) < 0$ and $v_p(\beta) = 0$, where $\overline{\mathbb{Q}}_p$ is an algebraic closure of the p-adic field $\mathbb{Q}_p$. This is possible because $\alpha\beta = \frac{1}{p}$ and we can take an automorphism of F and take an embedding of F into $\overline{\mathbb{Q}}_p$ such that $v_p(\alpha) < 0$ and $v_p(\beta) = 0$.

Since $b_0 \neq 0$ and $v_p(\beta) = 0$, the recurrence relation $(*)$ implies that for all large integers n, $v_p(c_n)$ is dominated by $v_p(\alpha^{n-1})$ which is negative. But for each N, there exists $n > N$ such that $c_n \neq 0$. Since a nonzero integer c_n has a nonnegative valuation, we get a contradiction. Hence, there is no such a sequence. $\square$

Proposition 2.8. *Let N be the conductor of the given elliptic curve $E/\mathbb{Q}$. If there is a nontrivial Heegner system attached to (E, K), where K is a quadratic imaginary extension of $\mathbb{Q}$ which is different from the field $\mathrm{End}(E) \otimes \mathbb{Q}$, then for a prime p such that $p \nmid r \cdot N \cdot [H_r : K] \cdot disc(H_r)$, there exist an integer r such that the rank of $E(H_{rp^n})$ is unbounded, as n goes to infinity.*

Proof. Let p be a prime $p \nmid r \cdot N \cdot [H_r : K] \cdot disc(H_r)$. As we have shown in Proposition 2.6, we show that for some integer r, the group generated by all of Heegner points of conductor rp^n for all integer $n \geq 1$ in the given nontrivial Heegner system is not finitely generated. Suppose for any integer r, this group is finitely generated. Then, since there are only finitely many torsion (Heegner) points over all the ring class fields in the system by Lemma 2.3, for some integer k, there is a fixed ring class field H_{rp^k} over which all Heegner points of conductor rp^n for all $n \geq 1$ are defined. Let $n_0 = rp^k$. Since the given Heegner system is nontrivial, we may assume that a Heegner point P_0 of conductor $n_0 = rp^k$ is of infinite order.

By the norm-compatibility property given in Proposition 2.2, we have that for all $n \geq 1$,

$$\mathrm{Trace}_{H_{n_0p^{n+1}}/H_{n_0p^n}}(P_{n+1}) = a_pP_n - P_{n-1},$$

where P_i are Heegner points of conductor n_0p^i.

Since P_{n+1} is defined over H_{n_0}, hence over $H_{n_0p^n}$ by assumption, the trace of P_{n+1} from $H_{n_0p^{n+1}}$ to $H_{n_0p^n}$ is the degree of $H_{n_0p^{n+1}}$ over $H_{n_0p^n}$ which is p by (a) of Lemma 2.4. Hence, the infinite sequence of Heegner points of conductor of the form n_0p^n satisfies the linear recurrence relation,

$$pP_{n+1} = a_pP_n - P_{n-1}, \text{ for all } n \geq 1. \tag{$**$}$$

By the Mordell-Weil Theorem, $E(H_{n_0})$ is finitely generated, so by dividing by its torsion subgroup $E(H_{n_0})_{tor}$, all points $P_n \mod E(H_{n_0})_{tor}$ lie in $\mathbb{Z}^k$ for some k. Suppose that $E(H_{n_0}) \cong \mathbb{Z}Q_1 + \cdots + \mathbb{Z}Q_k + E(H_{n_0})_{tor}$. Now we consider all points $P_n \mod E(H_{n_0})_{tor}$ and denote it by P_n again by abuse of natation. Let $P_n = \sum_{i=1}^{k} c_{n,i}Q_i$ for integers $c_{n,i}$. Since P_0 is not a torsion point by the assumption, without loss of generality we may assume that $c_{0,1} \neq 0$. Then, either $c_{2,1}$ or $c_{3,1}$ is nonzero, since otherwise, the relation $(**)$ implies that Q_1 is a linear combination of points $Q_2, Q_3, \ldots, Q_k$ over $\mathbb{Z}$ which contradicts the linear independence of points Q_i. By using $(**)$ and linear dependence of points Q_i inductively, we can show that $pc_{n+1} = a_pc_n - c_{n-1}$ for all $n \geq 1$, and if $c_{n,1} \neq 0$, then either $c_{n+1,1} \neq 0$ or $c_{n+2,1} \neq 0$. Hence, by letting $c_n = c_{n,1}$, we get an infinite sequence $\{c_n\}_{n=0}^{\infty}$ of integers satisfying the $pc_{n+1} = a_pc_n - c_{n-1}$ for all $n \geq 1$ with $c_0 \neq 0$ and for each N, there exists $n > N$ such that $c_n \neq 0$. This is impossible by Lemma 2.7. Therefore, we conclude that the rank of $E(H_{rp^n})$ is not finitely generated, as $n \to \infty$. In particular, by Lemma 2.5, we conclude that the rank of $E(H_{rp^n})$ cannot be bounded, as n goes to infinity. $\square$

Finally, the following proposition proves Theorem 1.4 hence, completes Theorem 1.3.

Proposition 2.9. *Let $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. Let K be a quadratic imaginary extension of $\mathbb{Q}$ such that $\sigma|_K \neq id_K$ and K is different from $\mathrm{End}(E) \otimes \mathbb{Q}$. Suppose all primes dividing the conductor N of E are split in K. Then, the rank of the Mordell-Weil group $E(H_n^\sigma)$ over the fixed subfield of H_n under σ is unbounded, as n goes to ∞. Hence, the rank of $E((K_{ab})^\sigma)$ is infinite, where K_{ab} is the maximal abelian extension of K.*

Proof. Since all primes dividing the conductor N of E are split in K, there is a nontrivial Heegner system attached to (E, K) by ([2], Chapter 3, Theorem 3.13). For a given $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, since $\sigma|_K \neq id_K$, the restriction of σ to each ring class field H_n in the given Heegner system can be lifted as an involution of H_n. Let $\sigma_n = \sigma|_{H_n}$ be the restriction of σ to H_n. Then, each ring class field H_n has a generalized dihedral group structure as its Galois group over $\mathbb{Q}$ with an involution σ_n such that for any $\tau \in \mathrm{Gal}(H_n/K)$, $\sigma_n \tau \sigma_n = \tau^{-1}$.

By Proposition 2.8, we fix a prime p and an integer r such that $p \nmid r \cdot N \cdot [H_r : K] \cdot disc(H_r)$ and the rank of $E(H_{rp^n})$ is unbounded, as n goes to infinity. We prove that for an odd prime p not dividing $rN[H_r : K]disc(H_r)$, the rank of $E(H^{\sigma}_{rp^n})$ is unbounded as n goes to infinity. Suppose not. Then since the restriction σ_n of σ to H_{rp^n} acts by an involution of each ring class field H_{rp^n}, there exists a fixed integer $n_0 = rp^k$ for some $k \geq 1$ such that σ acts by -1 on any nontrivial quotient $(E(H_{n_0p^n}) \otimes \mathbb{Q})/(E(H_{n_0}) \otimes \mathbb{Q})$, for all $n \geq 1$.

By (a) of Lemma 2.4, $\mathrm{Gal}(H_{n_0p^n}/H_{rp}) = \mathrm{Gal}(H_{rp^{k+n}}/H_{rp})$ is a cyclic group of order p^{k+n-1}. And $\mathrm{Gal}(H_{n_0p^n}/H_{n_0})$ is a subgroup of $\mathrm{Gal}(H_{n_0p^n}/H_{rp})$. Hence, it is a cyclic subgroup of order p^m for some $m < k + n - 1$.

Let τ_n be a generator of $\mathrm{Gal}(H_{n_0p^n}/H_{n_0})$. Consider $E(H_{n_0p^n}) \otimes \mathbb{Q}$ as a representation of $\mathrm{Gal}(H_{n_0p^n}/\mathbb{Q})$. And for each n, let

$$M_n = (E(H_{n_0p^n}) \otimes \mathbb{Q})/(E(H_{n_0}) \otimes \mathbb{Q}).$$

For every element $\alpha \in \mathrm{Gal}(H_{n_0p^n}/\mathbb{Q})$, let $\alpha|_{H_{n_0}}$ be the restriction of α to H_{n_0}. Then, $\alpha|_{H_{n_0}}$ is an element of $\mathrm{Gal}(H_{n_0}/K)$, since H_{n_0} is Galois over $\mathbb{Q}$. Therefore, $\mathrm{Gal}(H_{n_0p^n}/\mathbb{Q})$ acts on $E(H_{n_0}) \otimes \mathbb{Q}$ as well. So we can consider the quotient M_n as a representation of $\mathrm{Gal}(H_{n_0p^n}/\mathbb{Q})$.

Let

$$\rho : \mathrm{Gal}(H_{n_0p^n}/\mathbb{Q}) \to \mathrm{GL}(M_n)$$

be the representation of $\mathrm{Gal}(H_{n_0p^n}/\mathbb{Q})$. Then, by the hypothesis, σ_n acts by -1 on M_n. Hence, $\rho(\sigma_n) = -id$ on M_n. On the other hand, by the dihedral group structure of $\mathrm{Gal}(H_{n_0p^n}/\mathbb{Q})$, $\sigma_n \tau_n \sigma_n = \tau_n^{-1}$. Therefore,

$$\rho(\tau_n^2) = \rho(\tau_n)\rho(\tau_n) = (-id)\rho(\tau_n)(-id)\rho(\tau_n) = \rho(\sigma_n\tau_n\sigma_n\tau_n) = \rho(1) = id.$$

Hence, the restriction of ρ to the cyclic subgroup $\mathrm{Gal}(H_{n_0p^n}/H_{n_0})$ of $Gal(H_{n_0p^n}/\mathbb{Q})$ generated by τ_n^2 is a trivial representation of M_n. Since the order of τ_n is an odd integer p^m,

$$\langle \tau_n^2 \rangle = \langle \tau_n \rangle = \mathrm{Gal}(H_{n_0p^n}/H_{n_0}).$$

Therefore, we have

$$M_n^{\mathrm{Gal}(H_{n_0p^n}/H_{n_0})} = M_n^{\langle \tau_n^2 \rangle} = M_n.$$

This implies that

$$E(H_{n_0p^n}^{\mathrm{Gal}(H_{n_0p^n}/H_{n_0})}) \otimes \mathbb{Q} + E(H_{n_0}) \otimes \mathbb{Q} = E(H_{n_0p^n}) \otimes \mathbb{Q}.$$

Since $H_{n_0p^n}^{\mathrm{Gal}(H_{n_0p^n}/H_{n_0})} = H_{n_0}$,

$$E(H_{n_0}) \otimes \mathbb{Q} = E(H_{n_0p^n}) \otimes \mathbb{Q}, \quad \text{for all } n \geq 1,$$

which is a contradiction to Proposition 2.8.

Hence, the rank of $E(H_{n_0p^n}^\sigma)$ is unbounded, as $n \to \infty$. Since all ring class fields $H_{n_0p^n}$ are abelian over K, this implies that the rank of $E((K_{ab})^\sigma)$ is infinite. □

References

[1] C. Breuil, B. Conrad, F. Diamond, and R. Taylor, *On the modularity of elliptic curves over $\mathbb{Q}$: wild 3-adic exercises*, *J. Amer. Math. Soc.* **14** (2001), no. 4, 843-939.

[2] H. Darmon, *Rational points on modular elliptic curves*, CBMS Regional Conference Series in Mathematics, 101. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2004, (or see *http://www.math.mcgill.ca/darmon/pub/pub.html*).

[3] B. Im, *The rank of elliptic curves over large fields*, submitted for publication, 2003.

[4] M. Jarden and M. Fried, *Field Arithmetic*, A series of Modern Surveys in Math. **11**, Springer-Verlag, 1980.

[5] S. Lang, *Fundamentals of Diophantine Geometry*, Springer-Verlag, New York, 1983.

[6] M. Larsen, *Rank of elliptic curves over almost algebraically closed fields*, *Bull. London Math. Soc.* **35** (2003), 817–820.

[7] J.-P. Serre, *A course in Arithmetic*, Springer-Verlag, GTM **7**, 1973.

[8] J.-P. Serre, *Propriétés Galoisiennes des points d'ordre fini des courbes elliptiques*, *Invent. math.* **15** (1972), 259-331.

[9] J. H. Silverman, *Integer points on curves of genus* 1, *J. London Math. Soc.* **(2), 28** *(1983)*, 1-7.

[10] J. H. Silverman, *The Arithmetic of Elliptic Curves*, Springer-Verlag, GTM **106**, 1986.

[11] R. Taylor and A. Wiles, *Ring-Theoretic properties of certain Hecke algebras*, *Ann. of Math.* (2) **141** (1995), no. 3, 553-572.

[12] A. Wiles, *Modular elliptic curves and Fermat's last theorem*, *Ann. of Math.* (2) **141** (1995), no. 3, 443-551.

Department of Mathematics, Indiana University, Bloomington, Indiana 47405
E-mail address: `boim@indiana.edu`